

Green Geochemical Education in Libya: Current Status, Applications, Challenges, and Future Perspectives

And Tareq Alnnale³ Osama Rahil Shaltami¹, Mustafa A. Ben Hkoma²

osama.rahil@yahoo.com¹

¹Department of Earth Sciences, Faculty of Science, Benghazi University, Libya

²Libyan Centre for Sustainable Development Researches

Higher Institute of Science and Technology, Ragdalin – Libya³

t.alnaeli@histr.edu.ly³

<https://orcid.org/0009-0002-3282-4859>³

تاريخ الاستلام: 2026/04/01 تاريخ المراجعة 2026 /04/30 تاريخ القبول: 2026/05/13- تاريخ النشر: 2026 /06/16

Abstract

Dinosaur fossils are found in many Mesozoic rocks in Libya such as the Zarzaitine, Cabao, Mesak, Kiklah, and Mizdah formations. Dinosaur geochemistry in Libya is relevant to the fields of scientific investigation, geotourism, and sustainable development, as it contributes to the preservation and promotion of the country's fossil heritage. Although the relevance of dinosaur geochemistry is increasing worldwide, very few studies have been conducted regarding the geochemical aspects of dinosaurs from Libya. The scientific potential and possible avenues for future studies of dinosaur geochemistry in Libya are highlighted in this work.

Keywords: Dinosaur Geochemistry, Sustainable Development, Libya.

1. Introduction

The dinosaurs are a varied group of reptilian species that thrived on Earth during the Mesozoic Era. Dinosaurs first emerged some 230 million years ago in the Late Triassic and lasted until their extinction in the Late Cretaceous around 66 million years ago (Wicander, and Monroe, 2000; Erickson, 2002). The Triassic period was characterized by the dominance of small dinosaurs, while the Jurassic period was marked by the rapid diversification of large dinosaurs. The greatest diversity in dinosaurs was seen during the Cretaceous period; however, its end was marked by their extinction. Various factors led to the extinction of most non-avian dinosaurs, such as volcano eruption, meteorite collision, climatic change, and environmental disruption. Most of the dinosaurs were oviparous animals, including herbivores, carnivores, and omnivores. (Benton, 2019). Bones (e.g., Pyzalla et al., 2006; Britt et al., 2009; Kremer et al., 2012; An et al., 2016; Cerda et al., 2019; Garilli et al., 2023; Aureliano et al., 2026), teeth (e.g., Dyke and Malakhov 2004; Amiot et al., 2010; Montanari et al., 2013; Torices et al., 2018; Averianov et al., 2022; Michailow et al., 2025), footprints (e.g., Avanzini and Lockley, 2002; Lockley et al., 2006; Xing et al., 2011; Li et al., 2015; Liard et al., 2019; Antonelli et al., 2023; Jacobs, 2026), eggs (e.g., Codrea et al., 2002; Grellet-Tinner et al., 2006; Liang et al., 2009; Sellés et al., 2013; Tanaka et al., 2016; He et al., 2020; Xing et al., 2023; Kim et al., 2026), skin impressions (e.g., Kim et al., 2010; Rothschild and Depalma, 2013; Fondevilla et al., 2016; Hendrickx and Bell, 2021; Apesteuguía et al., 2023; Ralls, 2026), and coprolites—fossilized excrement— (e.g., Ghosh et al., 2003; Chin et al., 2009; Eriksson et al., 2011; Souto and

Fernandes, 2015; Mancuso et al., 2018; Dridi, 2022; Dino et al., 2026) are examples of dinosaur fossils. The major groups of dinosaurs are presented in Fig. 1.

Dinosaur geochemistry refers to the process by which geochemical techniques are used to analyze dinosaurs and the conditions under which they existed millions of years ago (e.g., Montanari et al., 2013; Vajda et al., 2016; Fanti et al., 2018; Ferrante et al., 2021). This field involves the use of geology, geochemistry, and paleontology to reconstruct past environments. The isotopes found in dinosaur fossils enable researchers to comprehend the depositional settings. Some of the most commonly used isotopes include those of carbon ($\delta^{13}\text{C}$), oxygen ($\delta^{18}\text{O}$), nitrogen ($\delta^{15}\text{N}$), and strontium ($^{87}\text{Sr}/^{86}\text{Sr}$). Major and trace elements in dinosaur fossils give information regarding their preservation, diagenesis, and depositional environment. The most important major elements are iron (Fe) and manganese (Mn), whereas rare earth elements (REE), uranium (U), and thorium (Th) are the significant trace elements. Biomarkers, proteins, lipids, and soft tissues are examined through the analysis of organic compounds found in fossils. Taphonomy provides insights into mineralization, bone preservation, fossil coloration, and burial settings. Geochemical analysis of rock strata containing dinosaur fossils is utilized in determining paleoclimatic conditions, ocean geochemistry, and volcanic eruptions. It aids researchers in understanding dinosaur evolution and extinction episodes. Conclusively, dinosaur geochemistry is an area that bridges paleontology and geochemistry and allows scientists to quantitatively establish how biological systems operated in the past using chemistry. This science forms a significant aspect of earth's evolution during the Mesozoic Era.

2. Objective

Reviewing the dinosaur geochemistry in Libya and connecting it to sustainable development is the goal of this study. In order to enhance knowledge of the geochemical importance of dinosaur fossils in Libya, the review entails the compilation and synthesis of preceding studies.




Fig. 1: Dinosaur classification simplified (after Kane and Willoughby, 2014).

3. Dinosaur Geochemistry in Libya

3.1. Dinosaur-Bearing Formations in Libya

The Cabao Formation (Early Cretaceous) which is located around Nalut (NW Libya) is the site of Libya's most significant dinosaur findings (Le Loeuff et al., 2010). There have been

reports of further dinosaur fossils from the Triassic Zarzaitine Formation (Hallett, 2002), the Early Cretaceous Mesak Formation (Abdullah et al., 2024), the Early Cretaceous Kiklah Formation (Smith and Vecchia, 2006) and the Late Cretaceous Mizdah Formation (Nessov et al., 1998). Fossils and replicas pertaining to Libyan dinosaur finds are preserved at the Nalut Dinosaur Museum (Fig. 2).

3.2. Sustainable Development Applications

3.2.1. Geotourism Development

One of the primary cornerstones of Libya's geoparks and sustainable tourism initiatives might be dinosaur fossils.

3.2.2. Scientific Research

Academic geosciences curricula are strengthened by dinosaur geochemistry. Fossil geochemistry research is becoming more and more important at Libyan universities.

3.2.3. Water and Environment

Depositional settings of dinosaur fossils may provide valuable insights into aquifers, paleoclimate, and desertification. This is consistent with the geosciences' role in sustainable development in Libya, particularly groundwater conservation and environmental surveillance.

3.2.4. Geoheritage Conservation

Many areas where dinosaur fossils have been found in Libya are still under-recorded, along with the illicit exploitation of such fossils. To achieve sustainable development, it is necessary that (1) There should be legal protection of the fossil-bearing sites; (2) Geoparks and museums should be established; (3) Digital documentation of the geosites; and (4) Awareness among local communities should be initiated.

References

- Abdullah, M.A., Albaghdady, A.A. and Abousif, A. (2024): Exploring newly discovered dinosaur trackways in the Messak Formation, Sebha region, southwest Libya. *Academy journal for Basic and Applied Sciences (AJBAS)*; 6: 1-11.
- Amiot, R., Kusuhashi, N., Xu, X. and Wang, Y. (2010): Isolated dinosaur teeth from the Lower Cretaceous Shapai and Fuxin formations of northeastern China. *Journal of Asian Earth Sciences*; 39(5): 347-358.
- An, W., Kuang, H., Liu, Y., Peng, N., Xu, K., Xu, H., Zhang, P., Wang, K., Chen, S. and Zhang, Y. (2016): Detrital zircon dating and tracing the provenance of dinosaur bone beds from the Late Cretaceous Wangshi Group in Zhucheng, Shandong, East China. *Journal of Palaeogeography*; 5(1): 72-99.
- Antonelli, M., Petti, F.M., Conti, J., Sacco, E., Petruzzelli, M., Spalluto, L. and Wagensommer, A. (2023): Lower Cretaceous dinosaur footprints from the Molfetta tracksite (Apulia, southern Italy). *Cretaceous Research*; 142: 105388.
- Apesteuguía, S., Díaz-Martínez, I., Mayoral, J.P., Riguetti, F., Veiga, G.D., de Valais, S. and Dumont, L.F. (2023): Dinosaur tracks with skin impressions in the La Buitrera Paleontological Area (Candeleros Formation, Cenomanian), Río Negro Province, Argentina. *Cretaceous Research*; 150: 105584.
- Aureliano, T., Maciel, V., Costa, V.P.G., de Paiva, A.C.F., de Aguilar Santos, C.L. and Ghilardi, A.M. (2026): Bone structure and the evolution of different pathways to gigantism in dinosaurs and megamammals. *Journal of South American Earth Sciences*; 169: 105855.
- Avanzini, M. and Lockley, M. (2002): Middle Triassic archosaur population structure: Interpretation based on *Isochirotherium delicatum* fossil footprints (Southern Alps, Italy). *Palaeogeography, Palaeoclimatology, Palaeoecology*; 185(3-4): 391-402.
- Averianov, A.O., Sizov, A.V., Grigoriev, D.V., Pestchevitskaya, E.B., Vitenko, D.D. and Skutschas, P.P. (2022): New data on dinosaurs from the Lower Cretaceous Murtoi Formation of Transbaikalia, Russia. *Cretaceous Research*; 138: 105287.
- Benton, M.J. (2019): *Dinosaurs rediscovered: The scientific revolution in paleontology*. Thames and Hudson Limited; 352p.
- Britt, B.B., Eberth, D.A., Scheetz, R.D., Greenhalgh, B.W. and Stadtman, K.L. (2009): Taphonomy of debris-flow hosted dinosaur bonebeds at Dalton Wells, Utah (Lower Cretaceous, Cedar Mountain Formation, USA). *Palaeogeography, Palaeoclimatology, Palaeoecology*; 280(1-2): 1-22.
- Cerda, I.A., Gasparini, Z., Coria, R.A., Salgado, L., Reguero, M., Ponce, D., Gonzalez, R., Jannello, J.M. and Moly, J. (2019): Paleobiological inferences for the Antarctic dinosaur *Antarctopelta oliveroi* (Ornithischia: Ankylosauria) based on bone histology of the holotype. *Cretaceous Research*; 103: 104171.
- Chin, K., Hartman, J.H. and Roth, B. (2009): Opportunistic exploitation of dinosaur dung: Fossil snails in coprolites from the Upper Cretaceous Two Medicine Formation of Montana. *Lethaia*; 42(2): 185-198.
- Codrea, V., Smith, T., Dica, P., Folie, A., Garcia, G., Godefroit, P. and Van Itterbeeck, J. (2002): Dinosaur egg nests, mammals and other vertebrates from a new Maastrichtian site of the Hațeg Basin (Romania). *Comptes Rendus Palevol*; 1(3): 173-180.
- Dino, M.A.P., de Andrade, R.C.L.P., Oliveira, N.C., Yamada, F.H., Saraiva, A.Á.F. and Bantim, R.A.M. (2026): Coprolites from the Romualdo Formation (Lower Cretaceous, Araripe Basin, Brazil): brief comments on morphology and trophic interactions. *Journal of South American Earth Sciences*; 171: 105931.
- Dridi, J. (2022): First report of diverse vertebrate coprofaunas from the Lower Cretaceous of Tunisia. *Cretaceous Research*; 135: 105192.

- Dyke, G.J. and Malakhov, D.V. (2004): Abundance and taphonomy of dinosaur teeth and other vertebrate remains from the Bostobynskaya Formation, north-east Aral Sea region, Republic of Kazakhstan. *Cretaceous Research*; 25(5): 669-674.
- Erickson, J. (2002): *Historical geology: Understanding our planet's past*. 2nd edition, Facts on File; 307p.
- Eriksson, M.E., Lindgren, J., Chin, K. and Månsby, U. (2011): Coprolite morphotypes from the Upper Cretaceous of Sweden: novel views on an ancient ecosystem and implications for coprolite taphonomy. *Lethaia*; 44(4): 455-468.
- Fanti, F., Bell, P.R., Tighe, M., Milan, L.A. and Dinelli, E. (2018): Geochemical fingerprinting as a tool for repatriating poached dinosaur fossils in Mongolia: A case study for the Nemegt Locality, Gobi Desert. *Palaeogeography, Palaeoclimatology, Palaeoecology*; 494: 51-64.
- Ferrante, C., Cavin, L., Vennemann, T. and Martini, R. (2021): Histology and geochemistry of *Allosaurus* (Dinosauria: Theropoda) from the Cleveland-Lloyd Dinosaur Quarry (Late Jurassic, Utah): Paleobiological implications. *Frontiers in Earth Science*; 9:641060.
- Fondevilla, V., Vila, B., Oms, O. and Galobart, À. (2016): Skin impressions of the last European dinosaurs. *Geological Magazine*; 154(2): 393-398.
- Garilli, V., Corrao, M., Grippi, S., Leotta, C., Sorbello, G., Galletti, L., Cillari, A., Guzzetta, D., Pollina, F., Sander, P.M. and Buffetaut, E. (2023): Ground Penetrating Radar to detect dinosaur bones within a Cretaceous hard limestone in Sicily. *Cretaceous Research*; 150: 105582.
- Ghosh, P., Bhattacharya, S.K., Sahni, A., Kar, R.K., Mohabey, D.M. and Ambwani, K. (2003): Dinosaur coprolites from the Late Cretaceous (Maastrichtian) Lameta Formation of India: isotopic and other markers suggesting a C₃plant diet. *Cretaceous Research*; 24(6): 743-750.
- Grellet-Tinner, G., Chiappe, L., Norell, M. and Bottjer, D. (2006): Dinosaur eggs and nesting behaviors: A paleobiological investigation. *Palaeogeography, Palaeoclimatology, Palaeoecology*; 232(2-4): 294-321.
- Hallett, D. (2002): *Petroleum geology of Libya*. Amsterdam, Elsevier Inc., 503p.
- He, Q., Yang, S., Jia, S., Xu, L., Xing, L., Gao, D., Liu, D., Gao, Y. and Zheng, Y. (2020): Trace element and isotope geochemistry of macroelongatoolithid eggs as an indicator of palaeoenvironmental reconstruction from the Late Cretaceous Xixia Basin, China. *Cretaceous Research*; 109: 104373.
- Hendrickx, C. and Bell, P.R. (2021): The scaly skin of the abelisaurid *Carnotaurus sastrei* (Theropoda: Ceratosauria) from the Upper Cretaceous of Patagonia. *Cretaceous Research*; 128: 104994.
- Jacobs, M.L. (2026): The historiography of dinosaur footprints of the Lower Cretaceous Wealden Group on the Isle of Wight, UK. *Proceedings of the Geologists' Association*; 137(1): 101164.
- Kane, J. and Willoughby, E. (2014): *Dinosaur classification simplified*. Emily Willoughby Art (available at <https://emilywilloughby.com/art/gallery/diagrams/dinosaur-classification-simplified>).
- Kim, J.Y., Kim, K.S., Lockley, M.G. and Seo, S.J. (2010): Dinosaur skin impressions from the Cretaceous of Korea: New insights into modes of preservation. *Palaeogeography, Palaeoclimatology, Palaeoecology*; 293(1-2): 167-174.
- Kim, S., Lee, Y., Kim, N. and Gihm, Y.S. (2026): Volcanic taphonomy of dinosaur eggs on Wi Island, South Korea: A reassessment of nesting chronology and cross-cutting relationships. *Cretaceous Research*; 183: 106347.
- Kremer, B., Owocki, K., Królikowska, A., Wrzosek, B. and Kazmierczak, J. (2012): Mineral microbial structures in a bone of the Late Cretaceous dinosaur *Sauroplophus angustirostris* from the Gobi Desert, Mongolia — a Raman spectroscopy study. *Palaeogeography, Palaeoclimatology, Palaeoecology*; 358-360: 51-61.

- Le Loeuff, J., Métais, E., Dutheil, D.B., Rubino, J.L., Buffetaut, E., Lafont, F., Cavin, L., Moreau, F., Tong, H., Blanpied, C. and Sbeta, A. (2010): An Early Cretaceous vertebrate assemblage from the Cabao Formation of NW Libya. *Geological Magazine*; 147(5): 750-759.
- Li, R., Lockley, M.G., Matsukawa, M. and Liu, M. (2015): Important Dinosaur-dominated footprint assemblages from the Lower Cretaceous Tianjialou Formation at the Houzuoshan Dinosaur Park, Junan County, Shandong Province, China. *Cretaceous Research*; 52(A): 83-100.
- Liang, X., Wen, S., Yang, D., Zhou, S. and Wu, S. (2009): Dinosaur eggs and dinosaur egg-bearing deposits (Upper Cretaceous) of Henan Province, China: Occurrences, palaeoenvironments, taphonomy and preservation. *Progress in Natural Science*; 19(11): 1587-1601.
- Liard, T., Liard, R. and Sutch, K. (2019): Dinosaur footprints from the Sao Khua Formation (Early Cretaceous) of Phu Kao National Park, Northeast Thailand. *Annales de Paléontologie*; 105(3): 217-222.
- Lockley, M.G., Houck, K., Yang, S., Matsukawa, M. and Lim, S. (2006): Dinosaur-dominated footprint assemblages from the Cretaceous Jindong Formation, Hallyo Haesang National Park area, Goseong County, South Korea: Evidence and implications. *Cretaceous Research*; 27(1): 70-101.
- Mancuso, A.C., Benavente, C.A., Previtera, E., Arcucci, A.B. and Irmis, R.B. (2018): Carnivore coprolites from the lower Carnian (Upper Triassic) Chañares Formation, northwestern Argentina. *Palaeogeography, Palaeoclimatology, Palaeoecology*; 489: 15-28.
- Michailow, M.M., Lugli, F., Cipriani, A., Giustina, F.D., Ferretti, A., Malferrari, D., Fowler, D., Fowler, E.F., Weber, M. and Tütken, T. (2025): Combined Ca, Sr isotope and trace element analyses of Late Cretaceous dinosaur teeth: assessing diet versus diagenesis. *Geochimica et Cosmochimica Acta*; 400: 172-189.
- Montanari, S., Higgins, P. and Norell, M.A. (2013): Dinosaur eggshell and tooth enamel geochemistry as an indicator of Mongolian Late Cretaceous palaeoenvironments. *Palaeogeography, Palaeoclimatology, Palaeoecology*; 370: 158-166.
- Nessov, A., Zhegallo, V.I. and Averianov, A.O. (1998): A new locality of Late Cretaceous snakes, mammals and other vertebrates in Africa (western Libya). *Annales de Paléontologie*; 84(3-4): 265-274.
- Pyzalla, A.R., Sander, P.M., Hansen, A., Ferreyro, R., Yi, S.-B., Stempniewicz, M. and Brokmeier, H.-G. (2006): Texture analyses of Sauropod dinosaur bones from Tendaguru. *Materials Science and Engineering: A*; 437(1): 2-9.
- Ralls, E. (2026): What colors were the gigantic, long-necked dinosaurs? Fossilized skin discovery tells us (available at <https://www.earth.com/news/fossilized-skin-discovery-diplodocus-sauropods-clues-about-dinosaur-color/>).
- Rothschild, B.M. and Depalma, R. (2013): Skin pathology in the Cretaceous: Evidence for probable failed predation in a dinosaur. *Cretaceous Research*; 42: 44-47.
- Sellés, A.G., Bravo, A.M., Delclòs, X., Colombo, F., Martí, X., Ortega-Blanco, J., Parellada, C. and Galobart, À. (2013): Dinosaur eggs in the Upper Cretaceous of the Coll de Nargó area, Lleida Province, south-central Pyrenees, Spain: Oodiversity, biostratigraphy and their implications. *Cretaceous Research*; 40: 10-20.
- Smith, J.B. and Vecchia, F.M.D. (2006): An abelisaurid (Dinosauria: Theropoda) tooth from the Lower Cretaceous Chicla formation of Libya. *Journal of African Earth Sciences*; 46(3): 240-244.
- Souto, P.R.F. and Fernandes, M.A. (2015): Fossilized excreta associated to dinosaurs in Brazil. *Journal of South American Earth Sciences*; 57: 32-38.
- Tanaka, K., Zelenitsky, D.K., Saegusa, H., Ikeda, T., DeBuhr, C.L. and Therrien, F. (2016): Dinosaur eggshell assemblage from Japan reveals unknown diversity of small theropods. *Cretaceous Research*; 57: 350-363.

- Temehu (2006): Nalut Dinosaur Museum (available at https://www.temehu.com/Cities_sites/museum-of-nalut.htm).
- Torices, A., Wilkinson, R., Arbour, V.M., Ruiz-Omeñaca, J.I. and Currie, P.J. (2018): Puncture-and-pull biomechanics in the teeth of predatory coelurosaurian dinosaurs. *Current Biology*; 28(9): 1467-1474.e2.
- Shaltami, O. R., & Hkoma, M. A. B. (2026). THE LINK BETWEEN DINOSAUR GEOCHEMISTRY AND SUSTAINABLE DEVELOPMENT IN LIBYA. *Al-Farooq Journal of Sciences*, 2(2), 229-238.
- Vajda, V., Fernández, M.D.P., Villanueva-Amadoz, U., Lehsten, V. and Alcalá, L. (2016): Dietary and environmental implications of Early Cretaceous predatory dinosaur coprolites from Teruel, Spain. *Palaeogeography, Palaeoclimatology, Palaeoecology*; 464: 134-142.
- Wicander, R. and Monroe, J.S. (2000): *Historical geology: Evolution of earth and life through time*. 3rd edition, Brooks Cole; 580p.
- Xing, L., Harris, J.D., Jia, C., Luo, Z., Wang, S. and An, J. (2011): Early cretaceous bird-dominated and dinosaur footprint assemblages from the northwestern margin of the Junggar Basin, Xinjiang, China. *Palaeoworld*; 20(4): 308-321.
- Alnnale, T. (2026). Predictive Governance in Digital Enterprises: An LSTM-Enhanced Deep Learning Framework for Economic Optimization of IT Incident Management Using Enriched Process Logs. *Al-Farooq Journal of Sciences*, 2(3), 86-113.
- Shaltami, O. R., Hkoma, M. A. B., Algomati, A. E., & Mohammed, A. A. F. K. (2026). War and Weapon Geochemistry: Key Areas, Applications and Impact on the Sustainable Development Goals. *Al-Farooq Journal of Sciences*, 2(2), 168-181.
- Azouz, A., Fawzi, M., Mohammed, I., Hamed, O., Maher, A., & Baddi, M. (2026). Influence of Electrolyte Chemistry and Electrode Material on Hydrogen Production Performance in Alkaline Water Electrolysis. *Al-Farooq Journal of Sciences*, 2(2), 49-66.
- Xing, L., Guo, Z., Wu, R., Du, C., Wang, D. and Persons IV, W.S. (2023): Cretaceous dinosaur and crocodylomorph egg records of the Sanshui Basin, Guangdong Province, southern China. *Journal of Palaeogeography*; 12(3): 401-418.