



Patient Reception and Registration in Outpatient Clinics, with a Focus on Patient Referrals from Clinics to Hospitals

Haneen Basheer Mohammed Al-Hamali¹, Maha Radwan Al-Jadidi², Haitham Mohammed Ahmad³

^{1,2,3}Faculty of Medical Technology, Misrata

Corresponding Author's Email: Haneen.alh@mtc.edu.ly

Received: 15/8/2026 Revised: 17/8/2026 Accepted: 05/9/2026 Publication: 25/10/2026

Abstract

This cross-sectional study examined 50 complete responses to a 25-item questionnaire on patient reception and registration, referral, coordination, electronic systems, and perceived quality and continuity of healthcare in Misrata, Libya. Composite scores were calculated as the mean of five items assigned to each dimension. Reception and registration had the highest mean ($M = 3.86$, $SD = 0.57$), followed by perceived quality and continuity ($M = 3.62$, $SD = 0.57$); electronic systems had the lowest mean ($M = 2.87$, $SD = 0.60$). Overall internal consistency was high (Cronbach's $\alpha = 0.987$). Pearson correlations ranged from $r = 0.916$ to $r = 0.956$, and all four were significant after Holm adjustment ($p < 0.001$); Spearman results were similar. Five groups of exact repeated response patterns involving 12 records were observed. A sensitivity analysis retaining one record per unique pattern yielded similar estimates ($n = 43$). The high reliability and correlations warrant cautious interpretation in view of potential item overlap and common-method effects; cross-sectional associations do not establish causation.

Keywords: patient reception; patient registration; referral system; outpatient clinics; electronic systems; healthcare quality; continuity of care.

1. Introduction

The organization of healthcare depends on the coordination of clinical work with the administrative processes that guide patients through a facility. Reception and registration are often the first points of contact. They help identify a patient, record essential information, locate an existing record, and direct the patient to an appropriate service. Errors or delays at this stage can complicate later communication and record retrieval.

The need for reliable information becomes more pronounced when a patient moves between a clinic and a hospital. A referral should communicate why further care is needed and provide relevant information to the receiving team. Referral systems also depend on clear responsibilities, appropriate destinations, and communication between providers. The World Health Organization's policy brief on referrals emphasizes the importance of defining services at each level and linking primary and secondary care as part of an organized system (World Health Organization, 2019).

Electronic systems may make referral information easier to prepare, transmit, and track. A scoping review found promising evidence that electronic referral systems can improve aspects of referral quality, communication, and workflow, while also noting that the evidence base is limited and varied across settings (Azamar-Alonso et al., 2019). Successful implementation is not only a technology issue: organizational arrangements, referral processes, coordination, and patient-related factors also matter (Seyed-Nezhad et al., 2021).

Against this background, the present study brings together reception and registration, clinic-to-hospital referral, inter-facility coordination, electronic systems, and perceived quality and continuity of healthcare. The analysis uses respondent-level data from 50 complete records to describe these dimensions, estimate internal consistency, and examine associations among composite scores.

1.1 Statement of the Problem

Healthcare institutions may face incomplete registration records, difficulty locating information, inconsistent referral documentation, delays, and limited communication between clinics and hospitals. These difficulties may make it harder for a receiving hospital to understand the reason for referral and may complicate subsequent follow-up. Paper-based or incompatible systems can add further barriers to timely information exchange. This questionnaire measures staff perceptions of these processes; it does not independently establish their operational prevalence or causes in the participating institutions.

The central research problem is therefore: What is the reported status of patient reception, registration, and clinic-to-hospital referral systems, and how are these processes associated with perceived quality and continuity of healthcare? The survey captures staff perceptions; it does not independently audit service efficiency or patient outcomes.

1.2 Research Questions

1. What is the reported level of efficiency of patient reception and registration?
2. What is the reported level of efficiency of patient referral from clinics to hospitals?
3. How do respondents assess coordination between clinics and hospitals?
4. What role do respondents attribute to electronic systems in registration and referral?
5. What relationships are reported between these administrative dimensions and perceived quality and continuity of healthcare?

1.3 Objectives of the Study

- Describe the reported status of patient reception and registration in outpatient services.
- Examine the reported status of referral procedures from clinics to hospitals.
- Assess respondents' reported views of coordination between referring and receiving facilities.
- Describe the reported use and perceived contribution of electronic systems.
- Describe item-level responses and compute dimension means, standard deviations, and internal-consistency estimates from respondent-level data.
- Develop practical recommendations for improving registration, referral documentation, coordination, and information exchange.

1.4 Significance of the Study

Scientifically, the study connects administrative processes that are sometimes examined separately: reception, registration, referral, coordination, and information systems. Practically, the comparatively lower ratings for coordination and electronic systems may help administrators identify areas for review. The findings are self-reported associations, not direct evidence of waiting times, referral completion rates, patient outcomes, or causal effects; operational indicators would be needed to assess those outcomes.

1.5 Hypotheses

The study tested four hypotheses: (1) patient reception and registration are positively associated with perceived quality and continuity of healthcare; (2) the patient referral system is positively associated with perceived quality and continuity; (3) coordination between clinics and hospitals is positively associated with perceived quality and continuity; and (4) electronic systems are positively associated with reception/referral efficiency. The fourth hypothesis was operationalized by correlating the Q16–Q20 electronic-systems composite with the mean of Q1–Q10. All four Pearson correlations were positive and statistically significant after Holm adjustment (all adjusted $p < 0.001$). These results are associations and do not demonstrate causal effects.

1.6 Scope of the Study

The study focuses on outpatient reception and registration and referral from clinics to hospitals, including coordination, electronic systems, and perceived quality and continuity. It analyzes 50 complete records from healthcare workers in Misrata.

2. Theoretical Framework and Previous Studies

2.1 Patient Reception, Registration, and Data Accuracy

Reception provides patients with initial guidance and directs them through the service. Registration records identifying and administrative information and can connect the current visit with an existing record. Accurate information supports retrieval, reduces avoidable duplication, and helps ensure that records are associated with the right person. These processes require consistent procedures for checking and correcting essential information, updating records, and protecting confidentiality. Electronic tools may assist these tasks, but they do not prevent errors if information is entered incorrectly or practices vary between staff.

2.2 Referral and Continuity of Care

A referral is a structured transition to another provider or level of care when further assessment or services are needed. A useful referral communicates the reason, urgency, relevant history, findings, and care already provided, while identifying the intended service. Continuity depends on more than sending a patient onward: information should accompany the patient, responsibilities should be clear, and appropriate follow-up or feedback should be possible. WHO guidance highlights the need to define services across levels and manage the link between primary and secondary care deliberately (World Health Organization, 2019).

2.3 Coordination and Electronic Referral

Coordination involves compatible referral procedures, accessible communication channels, clear ownership of incoming referrals, and a way to request missing information or communicate relevant outcomes. Electronic referral can support structured documentation and tracking. In their scoping review, Azamar-Alonso et al. (2019) found positive indications for referral quality and workflow, but emphasized that variation in studies limits generalization and that stronger evaluation is needed. Accordingly, electronic systems should be assessed in their local context rather than assumed to improve care simply because they are available.

A broader scoping review grouped factors affecting referral-system implementation into technological, process, organizational, and individual themes, including coordination and feedback, management and rules, and patient-related factors (Seyed-Nezhad et al., 2021). This supports a systems view: technology, staff training, policy, service capacity, and communication need to work together.

Empirical implementation studies of electronic referral provide relevant examples. Kim-Hwang et al. (2010) compared electronic and paper referrals at a public hospital, examining specialist assessments of referral information and process. Ulloa et al. (2017) evaluated a general-surgery electronic consultation system using a retrospective cohort of 150 referrals. Goh et al. (2025) used mixed methods and 25,404 referrals to examine electronic-record integration between primary and specialist care. Algerian et al. (2024) described referral patterns in Saudi Arabia's nationwide electronic-referral system, while Olson et al. (2021) evaluated a pilot of direct electronic referral and specialist access time.

Studies of communication, coordination, and continuity add patient and provider perspectives. O'Malley and Cunningham (2009) examined patient-reported coordination in relation to continuity and the primary-care physician's role as referral source. O'Malley and Reschovsky (2011) analyzed physician reports of communication during referrals and consultations. Zuchowski et al. (2015) documented referral-communication challenges and strategies in the Veterans Health Administration. Waibel et al. (2016) explored patients' perceptions of clinical and information continuity across care levels in Catalonia, and Juliani et al. (2017) examined Brazilian specialists'

perspectives on the referral process. These studies provide comparative context; their settings and measures differ from the present survey and do not validate its instrument or results.

2.4 Research Gap and Conceptual Focus

Prior literature addresses electronic referral and the organization of referral systems, but the practical conditions and resources of individual healthcare settings differ. This study focuses on the combined perceptions of healthcare workers regarding registration, referral, coordination, electronic tools, and quality/continuity in the reported Misrata setting. The individual records permit composite-score and correlation analyses, while the cross-sectional, self-reported design limits causal and external-validity conclusions.

3. Methodology

3.1 Research Design

The study used a descriptive-analytical, cross-sectional questionnaire design. Descriptive statistics, internal-consistency estimates, and prespecified associations between respondent-level composite scores were calculated from the individual records. No intervention or temporal follow-up is described, so causal effects cannot be established.

3.2 Population and Sample

The study sample comprised 50 complete responses from healthcare workers involved in reception, registration, referral, or related administrative procedures in Misrata. Of the respondents, 31 (62%) were women and 19 (38%) were men.

3.3 Instrument and Measurement

The questionnaire comprised 25 items scored on a five-point Likert scale from 1 (strongly disagree) to 5 (strongly agree). All 50 analyzed records contained valid integer responses from 1 to 5, with no missing item values. Items Q1–Q5 measured reception and registration, Q6–Q10 referral, Q11–Q15 coordination, Q16–Q20 electronic systems, and Q21–Q25 perceived quality and continuity.

3.4 Data Collection and Analysis

Analyses were conducted on the 50 complete respondent-level records. Item and composite means and sample standard deviations (denominator $n - 1$) were calculated. Internal consistency was assessed using Cronbach's alpha for each five-item dimension and for the full 25-item scale. Pearson product-moment correlations with two-tailed tests and Fisher-z 95% confidence intervals were used for the four prespecified associations; Holm adjustment was applied across the four Pearson tests. Spearman rank correlations were calculated as a sensitivity check for the ordinal Likert data. The fourth hypothesis was tested using the electronic-systems composite (Q16–Q20) and the combined reception/referral composite (Q1–Q10). A sensitivity analysis retaining one record per unique 25-item response pattern was also conducted ($n = 43$). No causal interpretation was made.

4. Results

4.1 Respondent Characteristics

The study included 50 records with unique respondent IDs and complete, valid item scores. A data-integrity audit identified five groups with exactly repeated response vectors across all 25 items, involving 12 records (seven repeats beyond one record per pattern). All 50 records were retained in the primary analysis. Table 1 summarizes the demographic characteristics of the respondents.

Characteristic	Category	Frequency	Percentage
Gender	Female	31	62%
Gender	Male	19	38%
Current workplace	Outpatient clinic	22	44%
Current workplace	Hospital	21	42%
Current workplace	Other healthcare facility	7	14%
Main work area	Reception and registration	12	24%
Main work area	Nursing	14	28%
Main work area	Medical practice	12	24%
Main work area	Medical records or administration	9	18%
Main work area	Other	3	6%
Work experience	Less than 1 year	7	14%
Work experience	1–5 years	21	42%
Work experience	6–10 years	13	26%
Work experience	More than 10 years	9	18%

Table 1. Respondent characteristics calculated from the 50 individual records.

4.2 Descriptive Results by Study Dimension

Table 2 reports composite means and sample standard deviations calculated directly from each respondent's five item scores in a dimension. Reception and registration had the highest mean ($M = 3.86$, $SD = 0.57$), while electronic systems had the lowest ($M = 2.87$, $SD = 0.60$). The overall 25-item mean was 3.32 ($SD = 0.59$).

Dimension	N	Items	Mean	SD	Cronbach's α
Patient reception and registration	50	5	3.86	0.57	0.923
Patient referral system	50	5	3.29	0.67	0.950
Coordination between clinics and hospitals	50	5	2.96	0.63	0.946
Electronic systems	50	5	2.87	0.60	0.944
Perceived quality and continuity of healthcare	50	5	3.62	0.57	0.937
Overall 25-item scale	50	25	3.32	0.59	0.987

Table 2. Composite scores are respondent-level means across five items; SD is the sample standard deviation. Cronbach's α was calculated from the raw items.

Table 3 and Figure 2 summarize the five response categories across the five items in each dimension (250 item responses per dimension). The response pattern shows a comparatively high proportion of “agree” responses for reception and registration, while neutral and disagree responses were more common in coordination and electronic systems.

Dimension	Strongly disagree (1)	Disagree (2)	Neutral (3)	Agree (4)	Strongly agree (5)
Patient reception and registration	1 (0.4%)	10 (4.0%)	50 (20.0%)	151 (60.4%)	38 (15.2%)
Patient referral system	2 (0.8%)	37 (14.8%)	105 (42.0%)	99 (39.6%)	7 (2.8%)
Coordination between clinics and hospitals	10 (4.0%)	53 (21.2%)	127 (50.8%)	58 (23.2%)	2 (0.8%)
Electronic systems	7 (2.8%)	67 (26.8%)	129 (51.6%)	45 (18.0%)	2 (0.8%)
Perceived quality and continuity of healthcare	1 (0.4%)	12 (4.8%)	91 (36.4%)	124 (49.6%)	22 (8.8%)

Table 3. Counts and percentages pooled across the five items in each dimension; denominator = 250 item responses per dimension.




Figure 1. Dimension composite means with 95% confidence intervals.




Figure 2. Five-point Likert response distribution pooled across items within each dimension.

4.2.1 Patient Reception and Registration

Item	Questionnaire statement	Mean	SD
Q1	Patients are received promptly upon arrival at the healthcare facility.	4.00	0.67
Q2	Registration procedures are clear and easy to follow.	3.78	0.62
Q3	Patient identification information is recorded accurately.	4.26	0.53
Q4	Patient records can be retrieved easily when needed.	3.26	0.83
Q5	Reception and registration staff provide patients with appropriate guidance.	4.00	0.57

Table 4.1. Item means and sample standard deviations calculated from the 50 individual responses.

4.2.2 Patient Referral System

Item	Questionnaire statement	Mean	SD
Q6	Referral procedures from clinics to hospitals are clearly defined.	3.16	0.84
Q7	Referral documents contain the information required by the receiving facility.	3.26	0.66
Q8	Patients are referred to appropriate hospitals or specialized services.	3.84	0.62
Q9	Referral procedures are completed within an appropriate timeframe.	3.00	0.73
Q10	Patients receive clear instructions regarding the referral process.	3.18	0.77

Table 4.2. Item means and sample standard deviations calculated from the 50 individual responses.

4.2.3 Coordination Between Clinics and Hospitals

Item	Questionnaire statement	Mean	SD
Q11	Communication between referring clinics and receiving hospitals is effective.	2.90	0.71
Q12	Responsibilities for managing referrals are clearly defined.	2.94	0.68
Q13	Hospitals can obtain additional information from referring clinics when needed.	3.62	0.64
Q14	Feedback about referred patients is communicated to the referring facility.	2.34	0.72
Q15	Coordination between clinics and hospitals supports continuity of patient care.	2.98	0.71

Table 4.3. Item means and sample standard deviations calculated from the 50 individual responses.

4.2.4 Electronic Systems

Item	Questionnaire statement	Mean	SD
Q16	Electronic systems are used effectively in patient registration and record management.	2.72	0.70
Q17	Electronic systems facilitate access to patient information when authorized.	3.02	0.77
Q18	Electronic tools improve the preparation and transmission of referral information.	2.78	0.68
Q19	Staff receive adequate training to use electronic registration and referral systems.	2.36	0.60
Q20	Appropriate measures protect the confidentiality of electronic patient information.	3.48	0.58

Table 4.4. Item means and sample standard deviations calculated from the 50 individual responses.

4.2.5 Perceived Quality and Continuity of Healthcare

Item	Questionnaire statement	Mean	SD
Q21	Reception and registration procedures contribute to the smooth delivery of healthcare services.	3.78	0.65
Q22	Accurate patient information helps healthcare professionals make appropriate care decisions.	4.14	0.57
Q23	Effective referral procedures help patients access the services they need.	3.74	0.63
Q24	Coordination between healthcare facilities supports follow-up after referral.	3.08	0.70
Q25	Administrative and referral procedures support the quality and continuity of healthcare.	3.34	0.63

Table 4.5. Item means and sample standard deviations calculated from the 50 individual responses.

4.3 Reliability and Inferential Analysis

Cronbach's alpha was high for the full 25-item scale ($\alpha = 0.987$) and ranged from 0.923 to 0.950 across the five dimensions (Table 2). These estimates indicate strong internal consistency in this sample, but do not establish construct validity. The very high full-scale alpha may also reflect item overlap and should be interpreted alongside the instrument's dimensional structure.

Table 5 reports the four prespecified Pearson correlations, with Fisher-z 95% confidence intervals and Holm-adjusted p-values. All associations were positive, very large, and statistically significant (all adjusted $p < 0.001$). Spearman rank correlations were similar, supporting the direction of the findings under a rank-based analysis. The fourth association uses Q16–Q20 for electronic systems and Q1–Q10 for reception/referral efficiency.

Hypothesized association	Pearson r	95% CI	Holm-adjusted p	Spearman p
Reception and registration vs perceived quality/continuity	0.956	[0.924, 0.975]	< .001	0.954
Referral system vs perceived quality/continuity	0.946	[0.907, 0.969]	< .001	0.955
Coordination vs perceived quality/continuity	0.938	[0.893, 0.965]	< .001	0.940
Electronic systems vs reception/referral efficiency (Q1–Q10)	0.916	[0.857, 0.952]	< .001	0.954

Table 5. Correlations between participant-level composite scores ($n = 50$); tests are two-tailed. Holm correction was applied to the four Pearson tests. All Holm-adjusted p-values were below .001.

The strength of these associations is unusually high for self-reported healthcare-process measures. The estimates should be interpreted cautiously because correlated measurement error, shared response tendencies, or overlapping item content may contribute to the observed associations. The cross-sectional design precludes causal interpretation.

The exact-response-pattern sensitivity analysis retained one record per distinct answer vector ($n = 43$). The overall mean was 3.34 and full-scale α was 0.986; dimension means differed from the primary analysis by no more than 0.033 points. Pearson correlations ranged from 0.932 to 0.952 and remained statistically significant (all $p < 0.001$). These estimates were similar to the primary analysis. The response matrix alone cannot determine whether identical patterns represent

independent responses or data-entry duplication; therefore, the reduced-record analysis is reported as a sensitivity analysis.

An exploratory item-level diagnostic found a mean Pearson inter-item correlation of 0.755 (range = 0.477–0.953). Under the five-item dimension assignment, the mean correlation among items within the same dimension was 0.766, compared with 0.753 between dimensions. Removing any single item yielded a full-scale alpha between 0.986 and 0.987. This pattern does not show clear separation among the five composites and may reflect overlapping item content or broad response tendencies. Given the sample size ($n = 50$), this descriptive check does not establish dimensional validity or common-method bias.

5. Discussion

The individual-level results show a clear gradient across dimensions. Reception and registration received the highest composite mean ($M = 3.86$, $SD = 0.57$), followed by perceived quality and continuity ($M = 3.62$, $SD = 0.57$) and referral ($M = 3.29$, $SD = 0.67$). Coordination ($M = 2.96$, $SD = 0.63$) and electronic systems ($M = 2.87$, $SD = 0.60$) received the lowest ratings. These scores describe the views of the 50 respondents and should not be treated as direct measures of service performance.

At item level, accurate recording of patient identification (Q3, $M = 4.26$, $SD = 0.53$) and accurate information for treatment decisions (Q22, $M = 4.14$, $SD = 0.57$) were rated most favorably. By contrast, communication of feedback to the referring facility (Q14, $M = 2.34$, $SD = 0.72$) and staff training in electronic systems (Q19, $M = 2.36$, $SD = 0.60$) had the lowest means. These findings suggest practical priorities for follow-up: strengthening the feedback loop after referrals and assessing staff training needs.

Referral to appropriate services (Q8, $M = 3.84$, $SD = 0.62$) was rated more favorably than completion within an appropriate timeframe (Q9, $M = 3.00$, $SD = 0.73$). Similarly, electronic confidentiality (Q20, $M = 3.48$, $SD = 0.58$) received a more favorable rating than effective system use (Q16, $M = 2.72$, $SD = 0.70$) or staff training (Q19, $M = 2.36$, $SD = 0.60$). These self-reports identify perceived strengths and constraints but cannot establish actual processing times, referral completion, interoperability, or security performance.

All four hypothesized associations were positive and very large: reception and registration with perceived quality/continuity ($r = .956$), referral with perceived quality/continuity ($r = .946$), coordination with perceived quality/continuity ($r = .938$), and electronic systems with the combined reception/referral score ($r = .916$); all Holm-adjusted p -values were below .001. Spearman results were similar. Such strong associations may reflect substantive relationships, overlapping item content, shared response tendencies, or common-method variance. The high full-scale alpha ($\alpha = .987$) reinforces the need to examine item redundancy and the intended five-factor structure rather than assuming that high internal consistency proves validity.

The results are consistent with literature emphasizing that referral systems depend on clear processes, information exchange, coordination, and staff capability (Azamar-Alonso et al., 2019; Seyed-Nezhad et al., 2021; World Health Organization, 2019). However, this study is cross-sectional and self-reported; neither the correlations nor the dimension means establish that a particular process causes better care or patient outcomes.

6. Conclusions

Analysis of the 50 complete records in the individual survey dataset found that reception and registration had the highest dimension mean ($M = 3.86$, $SD = 0.57$), while electronic systems had the lowest ($M = 2.87$, $SD = 0.60$). The overall 25-item mean was 3.32 ($SD = 0.59$), and Cronbach's alpha for the full scale was .987. The weakest item ratings concerned feedback to referring facilities and staff training in electronic systems.

The four hypothesized associations were positive and statistically significant after Holm adjustment, with Pearson correlations from .916 to .956. The magnitude of these associations should be

interpreted cautiously because they are unusually high, the measures are self-reported, and their constructs may overlap. Five groups of records shared exactly the same response vector across all 25 items, involving 12 records; a sensitivity analysis produced similar estimates when one record per unique pattern was retained. The findings support targeted review of referral feedback, staff training, and digital processes, but do not establish causal effects or objective service outcomes.

7. Recommendations

1. Standardize reception and registration procedures, including verification of essential identifiers, record updating, and correction of duplicate or inaccurate entries.
2. Use a referral form or structured referral record that captures the reason, urgency, destination, relevant clinical information, and contact details required for safe transfer.
3. Define who is responsible for sending, receiving, triaging, and following up each referral, with separate procedures for urgent cases.
4. Establish secure communication and feedback channels between clinics and hospitals, including a mechanism to request missing information and document referral outcomes.
5. Assess electronic systems against local infrastructure, interoperability, access control, training, maintenance, and continuity requirements before expanding their use.
6. Provide regular role-specific training on accurate data entry, referral documentation, confidentiality, and escalation procedures.
7. Monitor practical indicators such as incomplete referral forms, referral processing time, referral completion, and availability of feedback, while protecting patient confidentiality.
8. Preserve anonymized questionnaire data, the codebook, scoring rules, and statistical output in secure storage with an appropriate backup.

8. Limitations and Further Research

The cross-sectional self-report design limits causal inference and may be affected by common-method variance. Five groups with exact repeated response patterns involved 12 records; although the sensitivity estimates were similar, the response matrix alone cannot establish whether these patterns reflect independent responses or data-entry duplication. The sample size limits evaluation of the questionnaire's dimensional structure and generalizability. Findings should be interpreted as staff-reported perceptions rather than objective measures of service performance.

Further research should use a larger and clearly described sample, evaluate the instrument's dimensional structure, and test the stability of the observed correlations. Studies could combine staff perceptions with operational indicators such as referral completeness, processing time, completion, documented feedback, and patient outcomes. Any causal claims would require an appropriate longitudinal or intervention design.

References

- Aljerian, N. A., Alharbi, A. A., AlOmar, R. S., Binhotan, M. S., Alghamdi, H. A., Arafat, M. S., Aldhabib, A., & Alabdulaali, M. K. (2024). Showcasing the Saudi e-referral system experience: The epidemiology and pattern of referrals utilising nationwide secondary data. *Frontiers in Medicine*, *11*, Article 1348442. <https://doi.org/10.3389/fmed.2024.1348442>
- Azamar-Alonso, A., Costa, A. P., Huebner, L.-A., & Tarride, J. E. (2019). Electronic referral systems in health care: A scoping review. *ClinicoEconomics and Outcomes Research*, *11*, 325–333. <https://doi.org/10.2147/CEOR.S195597>
- Goh, K. H., Yeow, A. Y. K., Wang, L., Poh, H., Ng, H. J. H., Tan, G., Wee, S. K., Lim, E. L., & D'Souza, J. L. A. (2025). The benefits of integrating electronic medical record systems between primary and specialist care institutions: Mixed methods cohort study. *Journal of Medical Internet Research*, *27*, Article e49363. <https://doi.org/10.2196/49363>

- Juliani, C., MacPhee, M., & Spiri, W. (2017). Brazilian specialists' perspectives on the patient referral process. *Healthcare*, 5(1), Article 4. <https://doi.org/10.3390/healthcare5010004>
- Kim-Hwang, J. E., Chen, A. H., Bell, D. S., Guzman, D., Yee, H. F., Jr., & Kushel, M. B. (2010). Evaluating electronic referrals for specialty care at a public hospital. *Journal of General Internal Medicine*, 25(10), 1123–1128. <https://doi.org/10.1007/s11606-010-1402-1>
- O'Malley, A. S., & Cunningham, P. J. (2009). Patient experiences with coordination of care: The benefit of continuity and primary care physician as referral source. *Journal of General Internal Medicine*, 24(2), 170–177. <https://doi.org/10.1007/s11606-008-0885-5>
- O'Malley, A. S., & Reschovsky, J. D. (2011). Referral and consultation communication between primary care and specialist physicians: Finding common ground. *Archives of Internal Medicine*, 171(1), 56–65. <https://doi.org/10.1001/archinternmed.2010.480>
- Olson, E., Fruh, S., & Kleinpell, R. (2021). Electronic primary care to specialist referral: A pilot project to evaluate referral expedience and efficacy. *Journal of the American Association of Nurse Practitioners*, 33(4), 318–323. <https://doi.org/10.1097/JXX.0000000000000377>
- Seyed-Nezhad, M., Ahmadi, B., & Akbari-Sari, A. (2021). Factors affecting the successful implementation of the referral system: A scoping review. *Journal of Family Medicine and Primary Care*, 10(12), 4364–4375. https://doi.org/10.4103/jfmpc.jfmpc_514_21
- Ulloa, J. G., Russell, M. D., Chen, A. H., & Tuot, D. S. (2017). A cohort study of a general surgery electronic consultation system: Safety implications and impact on surgical yield. *BMC Health Services Research*, 17(1), Article 433. <https://doi.org/10.1186/s12913-017-2375-0>
- Waibel, S., Vargas, I., Aller, M.-B., Coderch, J., Farré, J., & Vázquez, M. L. (2016). Continuity of clinical management and information across care levels: Perceptions of users of different healthcare areas in the Catalan national health system. *BMC Health Services Research*, 16, Article 466. <https://doi.org/10.1186/s12913-016-1696-8>
- World Health Organization. (2019). *Strategies to strengthen referral from primary care to secondary care in low- and middle-income countries*. WHO Regional Office for South-East Asia. <https://apo.who.int/publications/9789290227090>
- Zuchowski, J. L., Rose, D. E., Hamilton, A. B., Stockdale, S. E., Meredith, L. S., Yano, E. M., Rubenstein, L. V., & Cordasco, K. M. (2015). Challenges in referral communication between VHA primary care and specialty care. *Journal of General Internal Medicine*, 30(3), 305–311. <https://doi.org/10.1007/s11606-014-3100-x>